

# Regional Distribution of Superoxide Dismutase Activity in Human Placenta and its Correlation with Lipid Peroxidation

Suleiman A. Saleh<sup>a,\*</sup>, Anwar D. Maraqa<sup>b</sup>, Mohammed Elamin Ali<sup>c</sup>, Zaki S. Mustafa<sup>c</sup>, Osama F. Qadoumi<sup>d</sup>

<sup>a</sup> Department of Laboratory Medical Sciences, Faculty of Allied Health Sciences, Hashemite University, Zarqa, Jordan .

<sup>b</sup> Faculty of Pharmacy and Medical Sciences, Al-Ahliyya Amman University, Amman, Jordan.

<sup>c</sup> Department of Obstetrics and gynaecology, College of Medicine, King Khalid University, Abha, Saudia Arabia.

<sup>d</sup> Department of Clinical Chemistry, College of Medicine, King Khalid University, Saudia Arabia. .

## Abstract

The objectives of the present study are to determine the activity of superoxide dismutase (SOD) in various regions of human placenta, and to correlate this activity with lipid peroxidation. To follow up changes in SOD activity and lipid peroxidation during gestation, a female rat model was used and pregnant females were sacrificed at 4, 7, 10, 14, and 18 days of gestation. Placentas were collected from 45 uncomplicated term pregnancies. Each placenta was sampled in 3 regions on the maternal and fetal side: central; mid placenta; and periphery. 5 gm of tissue were obtained for analysis and full thickness samples were obtained from the umbilical cord. In the rat, 2-3 placentas from each (female (n=4) were pooled and assayed while the cord for all placentas of the same female were pooled and assayed. The results show slight regional variations in the amount of lipid peroxidation as estimated by malondialdehyde (MDA) with the mid placental region of the fetal side being significantly higher than any of the other placental regions. MDA concentration in the placenta and umbilical cord of female rat progressively increased with progress of gestation (67% increase in the placenta and 90% increase in the umbilical cord). The results also show slight regional variations in SOD activity in the placenta being higher at the fetal site. The SOD activity in the umbilical cord was significantly higher than any of the placental regions. SOD activity in the placenta and umbilical cord of female rat progressively increased with the progress of gestation and after 10 days SOD activity in the umbilical cord was significantly higher than the placenta. The results clearly show that the increase of lipid peroxidation in the placenta and umbilical cord during gestation is coupled with an increase in SOD activity. The magnitude of SOD activity increase is higher to compensate for the increase in lipid peroxidation.

© 2010 Jordan Journal of Biological Sciences. All rights reserved

Keywords: placenta, umbilical cord, lipid peroxidation, SOD.

## 1. Introduction

Normal cells have a number of protective scavenger enzymes that act as antioxidants to detoxify the harmful reactive oxygen radical and prevent cell damage. One of these enzymes is superoxide dismutase (SOD: ECl.15.1.1) which catalyses the dismutation of superoxide radicals into oxygen and hydrogen peroxide and is widely distributed in mammalian tissues (McCord and Fridovich, 1969; Bannister *et al.*, 1987) including the placenta (Beckman, *et al.*, 1973; Sekiba and Yoshioka, 1979; and Takehara *et al.*, 1990). There are three SOD isoenzymes

(Beckman *et al.*, 1973) in human beings: the soluble cytosolic Cu/Zn SOD (Akbar *et al.*, 1998); the manganese-containing SOD (MnSOD) which is located in the matrix of the mitochondria (Bannister *et al.*, 1987), and the extracellular form of SOD (Marklund, 1984; Chu *et al.*, 2006).

Various studies dealt with the increase of lipid peroxidation and generation of free radicals during pregnancy (Ishihara 1978; Wickens *et al.*, 1981; Wang *et al.*, 1991; Walsh 1994; Akihito *et al.*, 2000; Warren *et al.*, 2005), but there is little information about placental lipid peroxides. It has also been suggested that placenta could contribute to the increase of lipid peroxidation (Diamant *et al.*, 1980; Hubel *et al.*, 1989), but little is known about the placental regional variation of this possible contribution.

Although the presence of Cu/Zn SOD activity in human placenta has been known for more than 30 years (Beckman *et al.*, 1973; Hien *et al.*, 1974) there are only sporadic

\* Corresponding author. S.Saleh@hu.edu.jo.

reports about the regional distribution of this enzyme in the placenta during gestation time. The objectives of this study are: (1) to measure the activity of the enzyme superoxide dismutase in various regions of human placenta immediately after delivery; (2) to determine the level of lipid peroxidation in the same placental regions; and (3) to correlate the level of lipid peroxidation with the activity of superoxide dismutase.

## 2. Materials and methods

### 2.1. Collection and processing of human placenta

Forty five placentas were collected from various maternity hospitals immediately after vaginal delivery and placed on ice. Each placenta was weighed and sampled in three main regions on both maternal and fetal sides as well as the umbilical cord as follows:

- a- central region: within 3 cm of the cord insertion,
  - b- mid placenta: mid way between the cord insertion and margin,
  - c- peripheral region: within 3 cm of the placental edge,
- umbilical cord: full thickness tissue samples were obtained from the umbilical cord (midway between the placenta and the umbilicus).

At each sampling site 5g of tissue was obtained, split into a liquotes and then kept in deep freeze (-80 C) until the time of processing.

### 2.2. Female rat model

Adult female rats weighing 250-300g were housed in groups of 4-5 in pvc cages (350x530 mm long x180 mm high) in an environment maintained at 19-22 C with 12/12 h light/ dark cycle. Food and water were always available. Female rats were allowed to mate with male rats and the males were removed after pregnancy which was confirmed with the observation of vaginal plug. Pregnant females were sacrificed at various gestational age (4, 7, 10, 14, and 18 days).

Embryos were carefully removed, placenta and umbilical cords were separated and processed similar to human placenta as mentioned in section 2.1. Each 2-3 placentas were pooled and assayed, but the cord of all placentas of the same female was pooled and assayed.

### 2.3. SOD assay

Placental tissues were homogenized in phosphate buffered saline (PBS, PH 7.4), sonicated and then centrifuged at 10.000 g for 25 min to obtain the cytosol fraction which was used to assay for Cu/Zn SOD. The assay was done according to the method of misra and Fridovich (1972) at 30 C . The amount of enzyme that causes 50% inhibition of epinephrine auto-oxidation is defined as 1 unit.

### 2.4. Measurement of lipid peroxidation

Lipid peroxidation was measured by determining the malondialdehyde (MDA) production using thiobarbituric acid (TBA) (Buege and Aust,1978). The MDA level was measured in the cytosol fraction of placental homogenate. Lipid peroxides were measured after addition of 2ml of TBA reagent (15% w/v trichloroacetic acid and 0.25 N

HCL) to 1 ml of cytosol fraction. The mixture was treated in a boiling water bath for 15 minutes. After cooling, the suspension was centrifuged at 1000 g for 10 minutes. The supernate was then separated and absorbance was measured at 535 nm. The MDA concentration was determined by the specific absorbance coefficient (1.34x10<sup>5</sup>mol/cm<sup>3</sup>).

### 2.5. Measurement of protein

Protein concentration in the cytosol suspension was determined by a modification of the Lowry procedure (1951) as reported by Markwell *et al.*, (1978) using bovine serum albumin as a standard.

### 2.6. Statistical methods

Statistical analysis was done using a t-test (SPSS for windows) by a personal computer. P values <0.05 were considered significant.

## 3. Results

### 3.1. Topographical data of maternal and placental records

The placental tissue was collected from 45 uncomplicated term pregnancies (38-40 weeks menstrual age). Mean weight of placenta was 720g (range 610- 945 g). Medical records for the women from whom placentas were recovered revealed that the mean maternal age was 28.4 years (range 19-38 years), mean gestational age was 38.6 weeks (range 37-42 weeks), mean parity was 3.7 (range 0-7), and mean birth weight of infants was 3028 g (range (2100-4030 g). Each placenta was sampled in 3 main regions on both the maternal and fetal sides. The various regions in the placenta which were dissected are indicated in Figure 1.

### 3.2. Regional distribution of SOD activity in the placenta

#### 3.2.1. Regional distribution of SOD in human placenta

Distribution of SOD activity in various regions of the placenta and the umbilical cord are shown in Table 1. There are slight regional variations in the SOD activity being relatively higher at the fetal site.

There is no significant difference in SOD activity of various regions at the same site of the placenta. However, the central region of the fetal site has a significantly higher level of SOD (P<0.05) than any of the regions in the maternal site. Activity of SOD was significantly higher (P < 0.01) in the umbilical cord than any of the regions in either the maternal or fetal sites of the placenta.

#### 3.2.2. SOD activity in placenta of female rat

Because the placenta of the rat embryo is much smaller than human placenta and the weight of each placenta was less than 0.5 g, regions of the placenta were not dissected. Two or three placentas were pooled and processed for SOD activity. Changes in the SOD activity, in the placenta




Diagram 1. Dissection of placental regions

Figure 1. Dissection of placental regions showing maternal and fetal sides of the placenta and the umbilical cord. Site of obtaining samples are indicated by \*. Modified after Jendryczko *et al.* 1991)

and the umbilical cord are shown in Table 2. SOD activity in either the placenta or the umbilical cord, progressively increased with the progression of gestational time. Until 7 days of gestation, there was no significant difference in SOD activity in the placenta compared to the umbilical cord. Although the activity of the enzyme in the placenta or the umbilical cord showed slight progressive increase, such increase was not significant. At 10 days of gestation, SOD activity in the placenta increased 58% while in the umbilical cord it increased about 2 folds. The increase in SOD activity in the placenta and the umbilical cord persisted throughout the rest of the gestational period. However, the increase was more evident in the umbilical cord. At 18 days, SOD activity in the placenta reached 2.1 folds, while in the umbilical cord it reached 3.1 folds.

### 3.3. Lipid peroxidation in the placenta

#### 3.3.1. Lipid peroxidation in human placenta

Lipid peroxidation in various placental regions is shown in Table 3. The average MDA concentration in the maternal side was 1.42 nmol/mg (range 1.41 – 1.53). In

spite of the slight regional variation, there was no significant difference ( $P > 0.05$ ). The average MDA concentration in the fetal side was 1.72 nmol/mg (range 1.62 – 1.83).

There are slight regional variations in the MDA concentration being relatively higher at the mid placental region. There is no significant difference in MDA concentration of various regions at the same site of the placenta.

However, the mid placental region of the fetal site was significantly higher ( $P < 0.05$ ) than any of the regions at the maternal site. Also the average MDA concentration in the fetal site was significantly higher ( $P < 0.05$ ) than the average maternal site. The MDA concentration in the umbilical cord was significantly higher than either the maternal or the fetal site of the placenta ( $P < 0.05$ ). It was 56% higher than the maternal site and 29% higher than the fetal site.

**Table 1: Regional distribution of SOD activity in human placenta and umbilical cord**

1. Placenta	
A. Maternal Side	
a. M1 (Central )	3.31 + 0.41
b. M2 (Mid placenta )	3.36 + 0.39
c. M3 (Periphery )	3.42 + 0.45
B. Fetal side	
a. F1 (central )	4.21 + 0.6*
b. F2 (Mid placenta )	4.13 + 0.53
c. F3 ( Periphery )	3.82 + 0.45
2. Umbilical cord	5.8 + 0.73**

$\bar{N} = 45$ , Values are mean + S.D , \*  $P < 0.05$  compared to maternal side ,  
\*\*  $P < 0.01$  Compared to placental region

**Table 2 : Changes of SOD activity in placenta and umbilical cord of female rat during gestation**

Gestation Age (days)	Fetus weight (g)	Placenta weight (g)	SOD activity ( unit/mg protein)	
			Placenta	Umbilical cord
4	1.4 + 0.2	0.23 + .02	2.6 + 0.3	2.5 + 0.4
7	2.1 + 0.2	0.31 + .03	2.7 + 0.4	3.1 + 0.5
10	3.1 + 0.3	0.35 + .03	4.1 + 0.4*	5.2 + .07*
14	3.6 + 0.4	0.38 + .04	4.6 + 0.5*	5.7 + .08*
18	4.4 + 0.5	0.45 + .05	5.5 + 0.6 **	7.8 + .09 **

$N = 4$  female rats, 2-3 placentas were pooled from each female rat and umbilical cords of all placentas of the same female were pooled. values are mean + S.D, \*  $P < 0.05$  , \*\*  $P < 0.01$

**Table 3. Lipid peroxidation in various regions of human placenta as estimated by MDA**

Type of Tissue	MDA concentration nmol/ mg protein X 10 <sup>-1</sup>
<b>1. Placenta</b>	
<b>A. Maternal side</b>	
a. M1 (central)	1.41 + 0.16
b. M2 (mid placental)	1.32 + 0.14
c. M3 (peripheral)	1.53 + 0.16
<b>B. Fetal side</b>	
a. F1 (central)	1.62 + 0.18
b. F2 ( mid placenta )	1.83 + 0.21*
c. F3 (peripheral)	1.71 + 0.19
<b>2. Umbilical cord</b>	<b>2.21 + 0.23**</b>

N=45, values are mean + S.D , \*P< 0.05 compared to maternal side.  
\*\*P<0.01 compared to placental region

### 3.3.2. Lipid peroxidation in the placenta of female rat

Lipid peroxidation in the placenta of female rat is shown in Table 4. MDA concentration in the placenta and the umbilical cord progressively increased with the progression of gestational time. Until 10 days of gestation, there was no significant difference in lipid peroxidation between the placenta and umbilical cord. At 14 and 18 days the difference in MDA concentration between the placenta and the umbilical cord was significant ( $P < 0.05$ ), with the umbilical cord being higher.

MDA concentration in the placenta increased 67% at 18 days while in the umbilical cord it increased 90%.

## 4. Discussion

The current study determined the level of lipid peroxidation in human and female rat placentas. We found that the level of peroxidation in both the human and the rat placenta are comparable. Concentration of peroxidation as estimated by MDA in the placenta of female rat at 18 days (2 days before delivery) was higher than that in full term delivered human placenta. The level was 2.45 nmol/mg in the rat versus 1.5 nmol/mg (average of the maternal and fetal regions) in humans (Table 3 and 4). The level is 63% higher in female rat.

The placenta is a heterogeneous tissue with membranous nature (Ali *et al.*, 1996); so different tissue compartments probably contributed to the tissue level of lipid peroxides (Roijsmakers *et al.*, 2004). This could explain the slight variation in the level of peroxidation at

various regions of the placenta being relatively higher at the fetal site. In both human and female rat, the level of lipid peroxidation in the umbilical cord was significantly higher than the placenta. In human placenta, the level of peroxidation in the umbilical cord was 47% higher than the placenta while in the female rat the level of peroxidation was about 12% higher. This indicates that the umbilical cord has a considerable contribution to the formation of lipid peroxides.

The level of peroxidation in the placenta or the umbilical cord, as shown in the present study, is high compared to other tissues like liver or lung as reported by Beckman *et al.*, 1973; Romero *et al.*, 1998; Janssen and Tazzerio, 2002). This high level could be due to increased thromboxane production as reported by various investigators (Wang *et al.*, 1992; Walsh and Wang, 1995; Cueto *et al.*, 1997), which causes an increase in cyclooxygenase activity resulting in increased oxygen radical formation. Lipid peroxides are formed when oxygen radicals interact with polyunsaturated fatty acids. Since the placenta is a rich source of unsaturated fatty acids, the increased cyclooxygenase activity associated with increased thromboxane production could be coupled with increased placental peroxidation.

In this study, we also shown that lipid peroxidation progressively increases as gestation progress in both the placenta and the umbilical cord. The increase was 67% in

**Table 4. Lipid peroxidation in the placenta and umbilical cord of female rat during gestation as estimated by MDA**

Type of Tissue	MDA concentration nmol/ mg protein X 10 <sup>-1</sup>
<b>1. Placenta</b>	
<b>A. Maternal side</b>	
a. M1 (central)	1.41 + 0.16
b. M2 (mid placental)	1.32 + 0.14
c. M3 (peripheral)	1.53 + 0.16
<b>B. Fetal side</b>	
a. F1 (central)	1.62 + 0.18
b. F2 (mid placenta)	1.83 + 0.21*
c. F3 (peripheral)	1.71 + 0.19
<b>2. Umbilical cord</b>	<b>2.21 + 0.23**</b>

N=45, values are mean + S.D , \*P< 0.05 compared to maternal side.  
\*\*P<0.01 compared to placental region

the placenta and almost doubled in the umbilical cord (Table 4). Although several investigators (Sekiba and

Yoshika, 1979; Wickens, 1981; Wang *et al.*, 1992; Wang and Rogers, 1997) have also shown increased level of lipid peroxides during pregnancy (review Walch, 1994) this was in maternal blood and not in actual placental tissue as we have demonstrated. It is also worth mentioning that only scarce reports (Palan *et al.*, 2001) are available about lipid peroxidation in the umbilical cord compared to the placenta.


We have shown in a previous study that the walls of blood vessels do have considerable amount of lipid peroxidation and this is much higher in the varicose veins (Wali *et al.*, 2002).

In this study, we have also shown that SOD activity in the placenta exhibited slight regional variations being higher at the fetal site of the placenta, particularly the central region (Table 1). The SOD activity was significantly higher in the umbilical cord than any of the placental regions of either the maternal or the fetal site. The level of SOD in the placenta of female rat at 18 days of gestation was 48% higher than in the human placenta (5.5 units/mg in the female rat versus 3.71 units/mg in human).

The level of SOD activity in the umbilical cord of female rat was 35% higher than that in humans.

There was a progressive increase of SOD activity with progress gestation where the increase was 2 folds in the placenta and 3 folds in the umbilical cord (Table 2).

Results in this study clearly show that the increase of lipid peroxidation in the placenta and the umbilical cord during gestation is coupled with an increase in SOD activity. The magnitude of the increase in SOD activity is higher than the increase of lipid peroxidation in female rat. Correlation of the increase in lipid peroxidation and the increase in SOD activity is shown in Figure 2. This indicates that although lipid peroxidation increased with the progress of gestation, the antioxidant SOD activity has also increased to compensate for the increase in lipid peroxidation.



**Figure 2. Correlation between the magnitude of increase of lipid peroxidation and SOD activity in the placenta and the umbilical cord of female rat with progress of gestation**

Various investigators have reported the increase of lipid peroxidation during pregnancy (Diamant *et al.*, 1980; Hubel *et al.*, 1989; Walsh, 1994) which seems to play an important factor in the pathogenesis of pre-eclampsia.

Recently, there has been a growing interest in using antioxidants to suppress lipid peroxidation in various tissues including the placenta (Chow, 2001; Raijmakers, *et al.*, 2004). In a previous study, we have shown that treatment with vitamin E considerably reduced lipid peroxidation in spermatozoa and improved sperm motility. Therefore, supplementation with vitamin E could protect against the increase of lipid peroxidation in the placenta.

## References

- Akbar SA., Nicholaides KH, and Brown PR. 1998. The localization of copper/zinc superoxide dismutase in human normal term placenta by an immunofluorescence technique. *J. Obs. Gynecol* 18(1): 44-46
- Ali KZM, Burton, G.J, Morad N, and Ali ME. 1996. Does hypercapillarization influence the branching pattern of terminal villi in the human placenta at high altitude. *Placenta* 17: 677-682
- Bannister JV, Bannister WH, and Rotilio, G. 1987. Aspects of the structure, function, and application of superoxide dismutase. *CRC Critical review Biochem* 22: 111-180
- Beckman G., Lundgren E., and Tarvik A. 1973. Superoxide dismutase isozymes in different human tissues, their genetic control and intracellular localization. *Human Heredity* 23: 338-345
- Buchanan M. 2006. Biological basis and clinical implications of acetylsalicylic acid resistance. *Can J Cardiol* 22: 149-151
- Chu y., Piper R., Yoshimasa W., Patel, P., and Heistad D.D. 2006. Endocytosis of extracellular superoxide dismutase into endothelial cells. *Atheros Thrombo Vas Biol.* 26: 1985-1990
- Cueto SM, Romany AD, Wang Y, and Walsh S. 1997. b- carotene attenuates peroxide-induced vasoconstriction in the human placenta. *J Soc Gynecol Invest* 4: 64-71
- Diamant S, Kissilewitz R, and Diamant Y. 1980. Lipid peroxidation system in human placental tissue: general properties and the influence of gestational age. *Biol Reprod* 23: 776-781
- Haliwell, B. 1996. Antioxidants in human health and disease. *Ann Rev Nut.* 16: 33-50
- Haliwell B, and Gutteridge JM. 1990. The antioxidants of human extracellular fluids. *Arc Biochem Biophys* 280: 1-8
- Hien PV, Kovacs K, and Matkovic B. 1974. Properties of enzymes, study of superoxide dismutase activity changes in human placenta of different ages. *Enzyme* 18: 341-347
- Hubel CA, Roberts I.M, Tatlor N, Musci, T.I., and Rogers G.M., McLaughlin M.K. 1989. Lipid peroxidation in pregnancy: New perspectives on preeclampsia. *Am. J. Obs Gynecol* 61: 1025-1034
- Ishihara M. 1978. Studies on lipoperoxide of normal pregnant women and of patients with toxemia of pregnancy. *Clinica Chemica Acta* 84: 1-9
- Janssen J. and Tazzerro T. 2002. Involvement of TP and EP receptors in vasoconstrictor responses in pulmonary vasculature. *J Pharmacol Exp Ther* 301: 1060-1066
- Jendryczko A, Drozd M, and Wojcik A. 1991. Regional variations in the activities of antioxidant enzymes in the term human placenta. *Zentrablatt Fur Gynakologie* 113 (19): 1059-1066
- Kulkarni AP and Kenel MF. 1987. Human placenta lipid peroxidation, some characteristics of NADPH microsomal fraction. *Gen Pharma* 18(5): 491-496
- Lowry OH, Rosenbrough NJ, Farr AL, Ranadall RG. 1951. Protein measurement with the folin phenol reagent. *J Biol Chem* 198: 265-268
- Marklund SL. 1984. Extracellular superoxide dismutase in human tissues and human cell lines. *J Clin Invest* 74: 1398-1403
- Markwell MAK, Haas SM, Bieber LL, and Tolbert NE. 1978. A modification of the Lowry procedure to simplify protein determination in membrane and lipoprotein samples. *Anal Biochem* 87: 206-210.
- McCord JM and Fridovich I. 1969. Superoxide dismutase. An enzyme function for erythrocyte (hemocuprein). *J Biol Chem* 244: 6049-6055
- Misra HP, and Fridovich I. 1962. The role of superoxide anion in the autooxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem* 247: 3170-3174
- Polan PR, Mikhail MS, Romney SL. 2001. Placental and serum levels of carotenoids in preeclampsia. *Obs Gynae* 98: 459-462
- Raijmakers MT, Decherd R, Poston L. 2004. Oxidative stress and preeclampsia: Rational for antioxidant clinical trials. *Hypertension* 44: 347-384.
- Romero JB, Bosch-Morell F, Romero MJ, and Enrique J. 1998. Lipid peroxidation products and antioxidants in human disease. *Health Prospect. Supp* 106(5): 20-24
- Sekiba K and Yoshioka T. 1979. Changes of lipid peroxides and superoxide dismutase activity in the human placenta. *Am J Obs Gynecol* 135: 369-371
- Suleiman SA, Ali EM, Zaki MS, Elmalik A, Nasr MA. 1996. Lipid peroxidation and human sperm motility: Protective role of vitamin E. *J Androl.* 17(5): 530-537
- Takehara Y, Yoshioka T, Sasaki J. 1990. Changes in the levels of lipoperoxide and antioxidant factors in human placenta during gestation. *Acta Med Okayama* 44(2): 103-111
- Uotila JT, Tuimala RJ, and Aarino TM. 1993. Findings on lipid peroxidation and antioxidant function in hypertensive complications of pregnancy. *J Obs Gynecol* 100: 270-276
- Walsh SW and Wang Y. 1993. Secretion of lipid peroxides by human placenta. *Am J Obs Gynecol* 169: 1462-1466
- Walsh SW. 1994. Lipid peroxidation in pregnancy. *Hyperten Pregnan* 13(1): 1-32.
- Walsh SW and Wang Y. 1995. Trophoblast and placental villous core production of lipid peroxides, thromboxanes and prostacyclin in pre-eclampsia.
- Wali MA, Suleiman SA., Kadoumi OF, and Nasr, M.A. 2002. Superoxide radical concentration and superoxide dismutase (SOD) enzyme activity in varicose veins. *Ann Thoracic cardiovas Surg* 8: 286-290

- Wang Y, Walsh SW, Guo J, and Zhang J. 1991. Maternal levels of prostacyclin, thromboxane, vitamin E, and lipid peroxides throughout normal pregnancy. *Am J Obs Gynecol* 165: 1690-1694
- Wang Y, Walsh SW, and Kay HH. 1992. Placental lipid peroxides and thromboxanes are increased and prostacyclin decreased in women with preeclampsia. *Am J Obs Gynecol*. 169:1462-1466
- Wang CC and Rogers MS. 1997. Lipid peroxidation in cord blood : effect of amniotic fluid volume. *Br J Obs Gynecol*. 104: 1140-1144

Copy Rights